SUMMARY OF WATER CONDITIONS May 1, 2002

Precipitation during April was disappointing, not quite half average for the month. About half the snowpack of a month ago has melted, an unusually high fraction for April. Because of the dryness of the spring so far, runoff forecasts have been reduced about 5 percent from those published on April 1. Although the outlook is not as bright as it was a month ago, runoff in most areas will be much better than last year except in the southern end of the State. Most users should have adequate supplies; however, significant delivery deficiencies are expected for Delta exporters and in local surface supplies in the south.

Forecasts of April through July runoff are now expected to be 75 percent of average statewide, but lower in the southern end of the Sierra. Water year forecasts are also about 75 percent of average compared to 50 percent actual runoff last year.

Snowpack water content is about 60 percent of average and about 45 percent of the April 1 average, the normal date of maximum accumulation. A number of sunny, warm days during the first part of April hastened the ripening process and produced much higher than normal early melt.

Precipitation during April was about 50 percent of average with a pronounced north to south gradient. Many basins in the southern portion of California had less than one quarter normal monthly precipitation, reinforcing the seasonal trend. Statewide precipitation this season since October 1 is about 80 percent of average compared to 75 percent last year.

Runoff so far this water year has been about 80 percent of average, almost double last year's 45 percent. April runoff overall was 85 percent of average, but a number of mountain rivers exceeded average runoff during the month due to accelerated snowmelt. Estimated runoff of the 8 major rivers of the Sacramento and San Joaquin River regions was 2.8 million acre—feet during April. The May estimate of the Sacramento River Index at the 90% exceedence level is 14.2 MAF. The May estimate of the San Joaquin Region 60–20–20 Index at the 75% exceedence level is 2.3.


Reservoir storage increased during April at an approximately average rate and is about average for the date. Except for the snowmelt driven river basins of the central and southern Sierra, storage has probably peaked for this season.

SUMMARY OF WATER CONDITIONS IN PERCENT OF AVERAGE

		—	• · · · · — · · ·			
HYDROLOGIC REGION	PRECIPITATION OCTOBER 1 TO DATE	MAY 1 SNOW WATER CONTENT	MAY 1 RESERVOIR STORAGE	RUNOFF OCTOBER 1 TO DATE	APR-JULY RUNOFF FORECAST	WATER YEAR RUNOFF FORECAST
NORTH COAST	95	65	100	85	85	85
SAN FRANCISCO BAY	100		90	90		
CENTRAL COAST	70		100	35		
SOUTH COAST	30		80	10		
SACRAMENTO RIVER	95	65	100	80	75	80
SAN JOAQUIN RIVER	85	65	105	75	75	70
TULARE LAKE	80	45	90	70	65	60
NORTH LAHONTAN	90	65	65	75	75	70
SOUTH LAHONTAN	35	20	110	80	75	80
COLORADO RIVER- DESERT	5					
STATEWIDE	80	60	100	80	75	75


DEPARTMENT OF WATER RESOURCES CALIFORNIA COOPERATIVE SNOW SURVEYS SEASONAL PRECIPITATION

IN PERCENT OF AVERAGE TO DATE
October 1, 2001 through April 30, 2002



DEPARTMENT OF WATER RESOURCES CALIFORNIA COOPERATIVE SNOW SURVEYS

FORECAST OF APRIL – JULY UNIMPAIRED SNOWMELT RUNOFF May 1, 2002



^{*} FORECAST BY DEPARTMENT OF WATER AND POWER, CITY OF LOS ANGELES

MAY 1, 2002 FORECASTS APRIL-JULY UNIMPAIRED RUNOFF

	Unimpaired Runoff in 1,000 Acre-Feet (1)								
HYDROLOGIC REGION	н	•					RECAST		
and Watershed	50 Yr	Max	Min	Apr-Jul	Pct	80 %	%		
	Avg	of	of	Forecasts	of	Probab			
	(2)	Record	Record		Avg	Range	-		
SACRAMENTO RIVER	()			1	<u> </u>				
Upper Sacramento River									
Sacramento River at Delta above Shasta Lake (3)	299	711	39	230	77%				
McCloud River above Shasta Lake	400	850	185	310	78%				
Pit River near Montgomery Creek + Squaw Creek	1,090	2,098	480	870	80%				
Total Inflow to Shasta Lake	1,849	3,525	726	1,450	78%	1,240 -	1,750		
Sacramento River above Bend Bridge, near Red Bluff	2,521	5,075	943	1,850	73%	1,570 -	2,290		
Feather River		.==	400	050	750/				
Feather River at Lake Almanor near Prattville (3)	333	675	120	250	75%				
North Fork at Pulga (3) Middle Fork near Clio (4)	1,028 86	2,416 518	243 4	730 55	71% 64%				
South Fork at Ponderosa Dam (3)	110	267	13	70	64%				
Feather River at Oroville	1,870	4,676	392	1,3 00	70%	1,080 -	1,600		
Yuba River	1,070	1,070	002	1,000	1070	1,000	1,000		
North Yuba below Goodyears Bar (3)	286	647	51	210	73%				
Inflow to Jackson Mdws and Bowman Reservoirs (3)	112	236	25	85	76%				
South Yuba at Langs Crossing (3)	233	481	57	170	73%				
Yuba River near Smartville plus Deer Creek	1,044	2,424	200	800	77%	690 -	940		
American River									
North Fork at North Fork Dam (3)	262	716	43	200	76%				
Middle Fork near Auburn (3)	522	1,406	100	410	79%				
Silver Creek Below Camino Diversion Dam (3)	173	386	37	130	75%				
American River below Folsom Lake	1,282	3,074	229	1,010	79%	880 -	1,180		
SAN JOAQUIN RIVER									
Cosumnes River at Michigan Bar	130	363	8	75	58%	45 -	105		
Mokelumne River									
North Fork near West Point (5)	437	829	104	340	78%				
Total Inflow to Pardee Reservoir	469	1,065	102	380	81%	340 -	450		
Stanislaus River									
Middle Fork below Beardsley Dam (3)	334	702	64	270	81%				
North Fork Inflow to McKays Point Dam (3)	224	503	34	180	80%	= 4.0	000		
Stanislaus River below Goodwin Reservoir (7)	716	1,710	116	580	81%	510 -	680		
Tuolumne River	222	707	07	240	750/				
Cherry Creek & Eleanor Creek near Hetch Hetchy (3) Tuolumme River near Hetch Hetchy (3)	322 606	727 1,392	97 153	240 470	75% 78%				
Tuolumne River below La Grange Reservoir (7)	1,230	2,682	301	940	76%	840 -	1,090		
Merced River	1,200	2,002	301	370	10/0		1,000		
Merced River at Pohono Bridge (3)	362	888	80	250	69%				
Merced River below Merced Falls (7)	633	1,587	123	420	66%	370 -	490		
San Joaquin River		,		_					
San Joaquin River at Mammoth Pool (6)	1,014	2,279	235	700	69%				
Big Creek below Huntington Lake (6)	95	264	11	65	68%				
South Fork near Florence Lake (6)	202	511	58	140	69%				
San Joaquin River below Millerton Lake	1,262	3,355	262	860	68%	760 -	980		
TULARE LAKE									
Kings River									
North Fork Kings River near Cliff Camp (3)	239	565	50	170	71%				
Kings River below Pine Flat Reservoir	1,234	3,113	274	860	70%	760 -	960		
Kaweah River below Terminus Reservoir	290	814	62	195	67%	165 -	235		
Tule River below Lake Success	65	259	2	28	43%	22 -	42		
Kern River									
Kern River near Kernville (3)	373	1,203	83	180	48%				
Kern River inflow to Lake Isabella	470	1,657	84	210	45%	180 -	270		

⁽¹⁾ See inside back cover for definition

⁽²⁾ All 50 year averages are based on years 1951-2000 unless otherwise noted

^{(3) 50} year average based on years 1941-90

^{(4) 44} year average based on years 1936-79

^{(5) 36} year average based on years 1936-72

^{(6) 45} year average based on years 1936-81

MAY 1, 2002 FORECASTS WATER YEAR UNIMPAIRED RUNOFF

				•		ired Ru								
Unimpaired Runoff in 1,000 Acre-Feet (1) HISTORICAL DISTRIBUTION FORECAST														
50 Yr	Max	Min	Oct	- .						Aug	Water	Pct	80 %	
Avg (2)	of Record	of Record	Thru Jan*	Feb *	Mar *	Apr *	May	Jun	Jul	& Sep	Year Forecasts	of Avg	Probab Range	-
(2)	Record	Record	Jan							Sep	rorecasis	Avg	Nange	(1)
888 1,234 3,217 6,194 8,990	1,965 2,353 5,150 10,796 17,180	165 557 1,484 2,479 3,294	2,275 3,835	520 810	575 830	480 650	420 520	300 380	250 300	450 545	5,270 7,870	85% 88%	5,000 - 7,510 -	5,610 8,360
780 2,417 219 291 4,775	1,269 4,400 637 562 9,492	366 666 24 32 994	1,090	305	445	505	450	220	125	160	3,300	69%	3,070 -	3,630
564 181 379 2,459	1,056 292 565 4,926	102 30 98 369	495	170	280	330	330	110	30	35	1,780	72%	1,660 -	1,930
616 1,070 318 2,830	1,234 2,575 705 6,382	66 144 59 349	470	220	355	425	420	140	25	15	2,070	73%	1,930 -	2,250
409	1,253	20	59	32	60	37	25	10	3	1	227	56%	195 -	260
626 774	1,009 1,800	197 129	85	40	65	125	175	70	10	5	575	74%	530 -	650
471	929	88												
1,196	2,952	155	145	55	100	215	240	105	20	10	890	74%	820 -	1,000
461 770 1,974	1,147 1,661 4,631	123 258 383	245	80	140	305	385	210	40	15	1,420	72%	1,320 -	1,590
461 1,014	1,020 2,787	92 150	105	35	60	150	180	70	20	10	630	62%	580 -	710
1,337 112 248 1,851	2,964 298 653 4,642	308 14 71 362	155	55	95	245	355	200	60	35	1,200	65%	1,090 -	1,340
284 1,736 460	607 4,287 1,402	58 386 94	145 60	50 17	80 30	235 62	360 85	205 38	60 10	35 8	1,170 310	67% 67%	1,060 - 270 -	1,280 360
153 558 741	615 1,577 2,318	163 175	30 75	6 20	30	13 55	10	4 50	1 25	0 25	75 360	49% 49%	65 - 325 -	90
	,													

^{*} Unimpaired runoff in prior months based on measured flows

⁽⁷⁾ Forecast point names based on USGS gage names. Stanislaus below Goodwin also known as inflow to New Melones, Tuolumne River below La Grange also known as inflow to Don Pedro, Merced River below Merced Falls also known as inflow to McClure.

MAY 1, 2002 FORECASTS APRIL-JULY UNIMPAIRED RUNOFF

IVDDOL GOLO DEGLOV	Unimpaired Runoff in 1,000 Acre-Feet (1)							
HYDROLOGIC REGION		ISTORICAL		FORECAST				
and Watershed	50 Yr	Max	Min	Apr-Jul	Pct			
	Avg	of	of	Forecasts	of			
	(2)	Record	Record		Avg			
NORTH COAST								
Trinity River								
Trinity River at Lewiston Lake	660	1,593	80	580	88			
Scott River								
Scott River near Fort Jones	200	400	30	170	85			
Klamath River								
Total inflow to Upper Klamath Lake (3)	340	531	229	235	69			
Truckee River Lake Tahoe to Farad accretions	272	713	52 0.2	210	77			
Lake Tahoe Rise (assuming gates closed, in feet)	1.4	5.4	0.2	0.9	63			
Carson River					_			
West Fork Carson River at Woodfords East Fork Carson River near Gardnerville	55 190	135 407	12 43	45 145	81 76			
East 1 of the Galloot 1 through Gallano 1 this	100	101	.0	0				
Walker River								
West Walker River below Little Walker, near Coleville	153	330	35	115	75			
East Walker River near Bridgeport	65	209	7	35	54			
SOUTH LAHONTAN								
Owens River								
Total tributary flow to Owens River (4)	235	579	96	171	73			


⁽¹⁾ See inside back cover for definition

⁽²⁾ All 50 year averages are based on years 1951-2000 unless otherwise noted

⁽³⁾ Forecast by U.S. Natural Resources Conservation Service and National Weather Service California-Nevada River Forecast Center, May through September forecast, 30 year average based on years 1971-2000.

⁽⁴⁾ Forecast by Department of Water and Power, City of Los Angeles, average based on years 1951-2000.

Water Content in % of April 1 Average



NORTH COAST REGION

SNOWPACK– First of the month measurements made at 10 snow courses indicate an area wide snow water equivalent of 17.3 inches. This is 45 percent of the seasonal April 1 average and 65% of the May 1 average. Last year at this time the pack was holding 20.8 inches of water.


Precipitation

October 1 to date in % of Average 250 225 200 175 150 125 100 75 50 25 0 Smith Klamath Trinity Eel Russian

PRECIPITATION – Seasonal precipitation (October 1 through the end of last month) on this area was 95 percent of normal. Precipitation last month was about 75 percent of the monthly average. Seasonal precipitation at this time last year stood at 55 percent of normal.

Reservoir Storage


Contents of major reservoirs in % of capacity



RESERVOIR STORAGE– First of the month storage in 7 reservoirs was 2.5 million acre–feet which is 100 percent of average. About 80 percent of available capacity was being used. Storage in these reservoirs at this time last year was 95 percent of average.


Runoff

October 1 to date in % of average




RUNOFF – Seasonal runoff of streams draining the area totaled 9.6 million acre–feet which is 85 percent of the average for this period. Last year, runoff for the same period was 35 percent of average.

Water Content in % of April 1 Average




October 1 to date in % of Average




Reservoir Storage

Contents of major reservoirs in % of capacity



Runoff

October 1 to date in % of average



SACRAMENTO RIVER REGION

SNOWPACK— First of the month measurements made at 46 snow courses indicate an area wide snow water equivalent of 17.4 inches. This is 45 percent of the seasonal April 1 average and 65 percent of the May 1 average. Last year at this time the pack was holding 12.4 inches of water.


PRECIPITATION – Seasonal precipitation (October 1 through the end of last month) on this area was 95 percent of normal. Precipitation last month was about 45 percent of the monthly average. Seasonal precipitation at this time last year stood at 70 percent of normal.

RESERVOIR STORAGE– First of the month storage in 43 reservoirs was 13.3 million acre–feet which is 100 percent of average. About 85 percent of available capacity was being used. Storage in these reservoirs at this time last year was 90 percent of average.


RUNOFF – Seasonal runoff of streams draining the area totaled 11.2 million acre–feet which is 80 percent of average for this period. Last year, runoff for the same period was 55 percent of average.

The Sacramento Region 40–30–30 Water Supply Index is forecast to be 6.5 assuming median meteorological conditions for the remainder of the year. This classifies the year as "dry" in the Sacramento Valley according to the State Water Resources Control Board.

Water Content in % of April 1 Average




October 1 to date in % of Average




Reservoir Storage

Contents of major reservoirs in % of capacity



Runoff

October 1 to date in % of average



SAN JOAQUIN RIVER AND TULARE LAKE REGIONS

SNOWPACK– First of the month measurements made at 52 **San Joaquin Region** snow courses indicate an area wide snow water equivalent of 18.2 inches. This is 50 percent of the seasonal (April 1) average and 65 percent of the May 1 average. Last year at this time the pack was holding 21.2 inches of water.

At the same time 29 **Tulare Lake Region** snow courses indicated a basin–wide snow water equivalent of 10.0 inches which is 35 percent of the average for April 1 and 45 percent of May 1. Last year at this time the basin was holding 10.9 inches of water.

PRECIPITATION – Seasonal precipitation

(October 1 through the end of last month) on the **San Joaquin Region** was 85 percent of normal. Precipitation last month was about 40 percent of the monthly average. Seasonal precipitation at this time last year stood at 85 percent of normal. Seasonal precipitation on the **Tulare Lake Region** was 80 percent of normal. Precipitation last month was about 35 percent of the monthly average. Seasonal precipitation at this time last year stood at 85 percent of normal.


RESERVOIR STORAGE– First of the month storage in 34 **San Joaquin Region** reservoirs was 8.1 million acre—feet which is 105 percent of average. About 70 percent of available capacity was being used. Storage in these reservoirs at this time last year was 115 percent of average.

First of the month storage in 6 **Tulare Lake Region** reservoirs was 955 thousand acre–feet which is 90 percent of average and about 45 percent of available capacity. Storage in these reservoirs at this time last year was 105 percent of average.


RUNOFF– Seasonal runoff of streams draining the San Joaquin Region totaled 2.7 million acre–feet which is 75 percent of average for this period. Last year, runoff for the same period was 55 percent of average. Seasonal runoff of streams draining the Tulare Lake Basin totaled 927 thousand acre–feet which is 70 percent of average for this period. Last year runoff for this same period was 60 percent of average.

The San Joaquin Region 60–20–20 Water Supply Index is forecast to be 2.4 assuming median meteorological conditions. This classifies the year as "dry" in the San Joaquin River Region according to the State Water Resources Control Board.

Water Content in % of April 1 Average




October 1 to date in % of Average




Reservoir Storage

Contents of major reservoirs in % of capacity



Runoff

October 1 to date in % of average



NORTH AND SOUTH LAHONTAN REGIONS

SNOWPACK– First of the month measurements made at 3 **North Lahontan Region** snow courses indicate an area wide snow water equivalent of 14.1 inches. This is 55 percent of the seasonal (April 1) average and 65 percent of the May 1 average. Last year at this time the pack was holding 8.0 inches of water. At the same time 2 **South Lahontan** snow courses indicated a basin–wide snow water equivalent of 2.5 inches which is 15 percent of the seasonal (April 1) average and 20 percent of the May 1 average. Last year at this time the basin was holding 11.7 inches of water.

PRECIPITATION – Seasonal precipitation (October 1 through the end of last month) on the **North Lahontan Region** was 90 percent of normal. Precipitation last month was about 115 percent of the monthly average. Seasonal precipitation at this time last year stood at 50 percent of normal. Seasonal precipitation on the **South Lahontan** was 35 percent of normal. Precipitation last month was about 25 percent of the monthly average. Seasonal precipitation at this time last year stood at 105 percent of normal.


RESERVOIR STORAGE– First of the month storage in 5 **North Lahontan** reservoirs was 415 thousand acre–feet which is 65 percent of average. About 40 percent of available capacity was being used. Storage in these reservoirs at this time last year was 100 percent of average. Lake Tahoe was 1.7 feet above its natural rim on May 1. First of the month storage in 8 **South Lahontan** reservoirs was 288 thousand acre–feet which is 110 percent of average and about 70 percent of available capacity. Storage in these reservoirs at this time last year was 115 percent of average.

RUNOFF– Seasonal runoff of streams draining the **North Lahontan Region** totaled 327 thousand acrefeet which is 75 percent of average for this period. Last year, runoff for the same period was 55 percent of average.

Seasonal runoff of the Owens River in the **South Lahontan** totaled 63 thousand acre–feet which is 80 percent of average for this period. Last year runoff for this same period was 70 percent of average.


Precipitation

October 1 to date in % of Average




Reservoir Storage

Contents of major reservoirs in % of capacity



Runoff

October 1 to date in % of average



SAN FRANCISCO BAY AND CENTRAL COAST REGIONS

PRECIPITATION – Seasonal precipitation (October 1 through the end of last month) on the **San Francisco Bay Region** was 105 percent of normal. Precipitation last month was about 30 percent of the monthly average. Seasonal precipitation at this time last year stood at 85 percent of normal.

Seasonal precipitation on the **Central Coast Region** was 70 percent of normal. Precipitation last month was about 15 percent of the monthly average. Seasonal precipitation at this time last year stood at 110 percent of normal.

RESERVOIR STORAGE– First of the month storage in 18 **San Francisco Bay Region** reservoirs was 494 thousand acre–feet which is 90 percent of average. About 70 percent of available capacity was being used. Storage in these reservoirs at this time last year was 100 percent of average.

First of the month storage in 6 **Central Coast Region** reservoirs was 712 thousand acre–feet which is 100 percent of average and about 75 percent of available capacity. Storage in these reservoirs at this time last year was 125 percent of average.

RUNOFF– Seasonal runoff of the Napa River in the **San Francisco Bay Region** totaled 65 thousand acre–feet which is 90 percent of average for this period. Last year, runoff for the same period was 45 percent of average.

Seasonal runoff of streams draining the **Central Coast Region** totaled 110 thousand acre–feet which is 35 percent of average for this period. Last year runoff for this same period was 70 percent of average.

SOUTH COAST AND COLORADO RIVER REGIONS

PRECIPITATION – October through April (seasonal) precipitation on the **South Coast Region** was 30 percent of normal. April precipitation was 15 percent of the monthly average. Seasonal precipitation at this time last year was 95 percent of normal. Seasonal precipitation on the **Colorado River–Desert Region** was 5 percent of normal. Precipitation during April was 5 percent of average. Seasonal precipitation at this time last year stood at 160 percent of average.

RESERVOIR STORAGE – May 1 storage in 29 major **South Coast Region** reservoirs was 1.3 million acrefeet or 80 percent of average. About 65 percent of available capacity was being used. Storage in these reservoirs at this time last year was 90 percent of average.

On May 1 combined storage in Lakes Powell, Mead, Mohave and Havasu was about 38 million acre–feet or about 90 percent of average. About 70 percent of available capacity was in use. Last year at this time, these reservoirs were storing 102 percent of average.

RUNOFF – Seasonal runoff from selected **South Coast Region** streams totaled 4 thousand acre–feet which is 10 percent of average. Seasonal runoff from these streams last year was 35 percent of average.

COLORADO RIVER

The April July inflow to Lake Powell is forecast to be 2 million acre–feet, which is 25 percent of average. The May 1 snowpack in the Colorado River basin above Lake Powell was 25 percent of average, highest in the Upper Green at 70 percent and lowest in the Animas at below 5 percent.

CENTRAL VALLEY PROJECT

As of April 30, 2002 CVP storage was 9.6 million acre-feet which is an increase of 0.1 million acre feet compared to one year ago, and is approximately 113% of normal for that date.

The Bureau of Reclamation announced updated water allocations for the CVP contractors on April 15, 2002. Based on a conservative water supply forecast prepared from information available April 1, 2002, and a water year inflow into Shasta Reservoir of 5.0 million acre–feet, CVP water allocations were: Agricultural contractors North of Delta 100% and South of Delta 55%; Urban contractors North of Delta 100% and South of Delta 80%; Sacramento River water rights and San Joaquin Exchange Contractors 100%; Wildlife Refuges 100%; Friant Contractors will be 100 percent of Class 1 and 0 (zero) percent of Class 2. Updated allocations will be announced in mid–May. The forecast of CVP operations is available on the Mid–Pacific Region's website at www.mp.usbr.gov.

STATE WATER PROJECT

Total storage in the major SWP reservoirs was about 4.26 MAF on April 30, 2002, compared with 3.77 MAF at this time in 2001. On April 30 storage at Lake Oroville was about 2.66 MAF as compared to about 2.19 MAF last year.

The State's share of San Luis Reservoir storage at the end of April was 984 TAF, as compared to about 985 TAF at this time last year. This year's storage included approximately 42 TAF of water acquired byt the Environmental Water Account.

The combined storage of SWP's southern reservoirs was about 617 TAF on April 30, as compared to 604 TAF at this time last year.

SWP water deliveries through April 2002 were about 730 TAF. This is a combination of project, transfer, and exchange waters. This was about 194 TAF more than the same time period in 2001.

The SWP allocation was increased on March 28 to 2.5 MAF, which represents 60% of State Water Contractor entitlement.

MAJOR WATER DISTRIBUTION PROJECTS RESERVOIR STORAGE

(AVERAGES BASED ON 1951-2000 OR PERIOD RECORD)

RESERVOIR	CAPACITY 1,000 AF	AVERAGE STORAGE 1,000 AF	2001 1,000 AF	2002	RAGE AT EN PERCENT AVERAGE	PERCENT			
STATE WATER PROJEC		0.007	0.400	0.050	200/	750/			
Lake Oroville	3,538	2,967	2,188	2,659	90%	75%			
San Luis Reservoir (SWF	•	983	985	973	99%	92%			
Lake Del Valle	77	39	38	35	89%	45%			
Lake Silverwood	73	68	70	73	107%	100%			
Pyramid Lake	171	163	164	163	100%	95%			
Castaic Lake	324	286	258	263	92%	81%			
Perris Lake	132	117	112	117	100%	89%			
CENTRAL VALLEY PROJECT									
Trinity Lake	2,448	2,045	1,963	2,049	100%	84%			
Lake Shasta	4,552	3,950	4,020	4,297	109%	94%			
Whiskeytown Lake	241	231	236	240	104%	99%			
Folsom Lake	977	728	668	759	104%	78%			
New Melones Reservoir	2,420	1,446	1,876	1,589	110%	66%			
Millerton Lake	520	352	450	470	134%	90%			
San Luis Reservoir (CVP		880	926	871	99%	90%			
COLORADO RIVER PRO									
Lake Mead	26,159	20,374	21,603	18,539	91%	71%			
Lake Powell	25,002	19,267	18,821	16,705	87%	67%			
Lake Mohave	1,810	1,672	1,734	1,682	101%	93%			
Lake Havasu	619	588	594	584	99%	94%			
EAST BAY MUNICIPAL U	JTILITY DISTF	RICT							
Pardee Res	198	182	194	174	96%	88%			
Camanche Reservoir	417	258	270	296	114%	71%			
East Bay (4 reservoirs)	147	136	136	138	102%	94%			
CITY AND COUNTY OF	SAN FRANCIS	SCO							
Hetch-Hetchy Reservoir	360	157	217	180	114%	50%			
Cherry Lake	268	145	183	224	155%	84%			
Lake Eleanor	26	15	17	22	149%	84%			
Souty Bay/Peninsula (4 re	es.) 225	182	194	149	82%	66%			
CITY OF LOS ANGELES	S (D.W.P.)								
Lake Crowley	183	124	146	139	112%	76%			
Grant Lake	48	26	38	32	123%	67%			
Other Aqueduct Storage	(6 res.) 95	75	64	67	89%	71%			

TELEMETERED SNOW WATER EQUIVALENTS May 1, 2002 (AVERAGES BASED ON PERIOD RECORD)


INCHES OF WATER EC	QUIVALENT
--------------------	-----------

					R EQUIVALENT	
BASIN NAME		APRIL 1	F	PERCENT	24 HRS	1 WEEK
STATION NAME	ELEV	AVERAGE	May 1 OF A	VERAGE	PREVIOUS	PREVIOUS
TRINITY RIVER	LLL V	WEIGHOL	way i Oi i	WEIGIGE	TREVIOUS	TREVIOUS
Peterson Flat	7150'	29.2	16.4	56.1	16.4	19.3
			10.4	36.1	10.4	19.3
Red Rock Mountain	6700'	39.6	47.0	40.0	47.5	-
Bonanza King	6450'	40.5	17.8	43.9	17.5	21.6
Shimmy Lake	6400'	40.3	_	_	_	_
Middle Boulder 3	6200'	28.3	12.0	42.5	11.4	12.0
Highland Lakes	6030'	29.9	7.4	24.9	7.1	10.2
Scott Mountain	5900'	16.0	0.0	0.0	0.0	1.9
Mumbo Basin	5650'	22.4	_	_	_	_
Big Flat	5100'	15.8	3.4	21.6	3.4	7.0
SACRAMENTO RIVER	0.00	10.0	0.1	21.0	0.1	7.0
Cedar Pass	7100'	18.1	12.3	68.0	11.7	12.7
		12.7	12.3			12.7
Blacks Mountain	7050'		_		_	_
Sand Flat	6750'	42.4	20.3	47.8	19.6	22.0
Medicine Lake	6700'	32.6	16.9	51.9	16.4	18.7
Adin Mountain	6200'	13.6	1.1	8.1	0.6	2.5
Snow Mountain	5950'	27.0	12.0	44.4	12.0	15.7
Slate Creek	5700'	29.0	8.8	30.2	8.4	9.4
Stouts Meadow	5400'	36.0	4.4	12.3	4.4	6.4
FEATHER RIVER						
Kettle Rock	7300'	25.5	10.9	42.8	10.9	12.4
Grizzly Ridge	6900'	29.7	9.7	32.7	9.8	12.5
Pilot Peak	6800'	52.6	8.8	16.7	9.0	11.4
Gold Lake	6750'	36.5	33.5	91.7	33.5	34.2
Humbug	6500'	28.0	21.9	78.1	21.9	24.6
Rattlesnake	6100'	14.0	0.0	0.0	0.0	0.0
Bucks Lake	5750'	44.7	28.6	63.9	28.6	31.9
Four Trees	5150'	20.0	1.1	5.4	1.1	0.7
EEL RIVER						
Noel Spring	5100'	_	1.0	_	1.2	0.0
YUBA & AMERICAN RIVERS						
Lake Lois	8600'	39.5	_			_
			41.0	440.0	40.6	40.2
Schneiders	8750'	34.5		118.8		_
Caples Lake	8000'	30.9	17.4	56.3	17.3	18.6
Alpha	7600'	35.9	19.7	54.8	19.5	21.4
Meadow Lake	7200'	55.5	50.6	91.2	50.5	51.1
Silver Lake	7100'	22.7	7.3	32.2	7.1	10.2
Central Sierra Snow Lab	6900'	33.6	7.8	23.2	8.0	11.9
Huysink	6600'	42.6	29.9	70.1	29.6	31.1
Van Vleck	6700'	35.9	28.1	78.1	27.5	28.6
Robbs Saddle	5900'	21.4	10.1	47.1	9.9	12.0
Greek Store	5600'	21.0		-77.1	— —	12.0
	5280'	9.0	_	_	_	_
Blue Canyon			_	_	_	_
Robbs Powerhouse	5150'	5.2	0.0	0.0	0.0	0.0
MOKELUMNE & STANISLAUS RIV	-					
Deadman Creek	9250'	37.2	21.8	58.5	21.5	21.9
Highland Meadow	8700'	47.9	36.8	76.8	36.7	34.0
Gianelli Meadow	8400'	55.5	31.9	57.5	31.4	30.0
Lower Relief Valley	8100'	41.2	28.6	69.3	27.3	27.9
Blue Lakes	8000'	33.1	23.9	72.2	23.6	24.0
Mud Lake	7900'	44.9	43.8	97.5	43.5	43.6
Stanislaus Meadow	7750'	47.5	36.5	76.8	36.0	36.0
Bloods Creek	7200'	35.5	17.4	49.0	17.2	19.6
Black Springs	6500'	32.0	18.6	58.1	18.3	20.7
TUOLUMNE & MERCED RIVERS						
Tioga Pass Entrance	9945'	_	_	_	_	_
Dana Meadows	9800'	27.7	20.1	72.7	20.2	19.9
Slide Canyon	9200'	41.1	34.0	82.8	34.0	32.1
Lake Tenaya	8150'	33.1	20.7	62.5	20.0	19.4
Tuolumne Meadows	8600'	22.6	0.5	2.2	0.5	4.3
Horse Meadow	8400'	48.6	34.0	70.0	33.4	30.8
Ostrander Lake						
	8200'	34.8	19.6	56.3	18.9	19.6
Paradise Meadow	7650'	41.3	29.9	72.3	29.2	30.5
Gin Flat	7050'	34.2	11.5	33.7	11.3	13.3
Lower Kibbie Ridge	6700'	27.4	4.5	16.2	4.5	5.8

DACINIANAE		ADDII 4	IIN	DEDOENT		4 \\/\		
BASIN NAME		APRIL 1		PERCENT	24 HRS	1 WEEK		
STATION NAME	ELEV	AVERAGE	May 1 C	F AVERAGE	PREVIOUS	PREVIOUS		
SAN JOAQUIN RIVER	40050	20.4	04.0	00.0	00.5	22.0		
Volcanic Knob	10050'	30.1	24.2	80.3	23.5	22.9		
Agnew Pass Kaiser Point	9450' 9200'	32.3 37.8	10.8 9.2	33.3 24.3	10.1 9.2	11.4 10.8		
Green Mountain	7900'	30.8	9.2	24.3	9.2	10.0		
Tamarack Summit	7550'	30.5	1.4	4.5	1.4	4.5		
Chilkoot Meadow	7150'	38.0	16.1	42.3	16.0	17.5		
Huntington Lake	7000'	20.1	1.7	8.4	1.7	4.3		
Graveyard Meadow	6900'	18.8	0.0	0.0	0.0	2.5		
Poison Ridge	6900'	28.9	0.0	0.0	0.0	1.1		
KINGS RIVER								
Bishop Pass	11200'	34.0	18.7	55.0	18.7	20.7		
Charlotte Lake	10400'	27.5	21.0	76.4	21.6	22.9		
State Lakes	10300'	29.0	19.2	66.2	19.2	20.0		
Mitchell Meadow	9900'	32.9	27.5	83.6	27.2	27.4		
Blackcap Basin	10300'	34.3	25.7	75.0	25.7	25.4		
Upper Burnt Corral	9700'	34.6	25.3	73.2	25.3	25.3		
West Woodchuck Meadow	9100'	32.8	15.6	47.6	15.0	16.3		
Big Meadows	7600'	25.9	0.8	3.1	1.3	5.3		
KAWEAH & TULE RIVERS								
Farewell Gap	9500'	34.5	22.5	65.2	22.5	27.1		
Quaking Aspen	7200'	21.0	0.0	0.0	0.0	0.0		
Giant Forest	6650'	10.0	0.0	0.0	0.0	0.0		
KERN RIVER Upper Tyndall Creek	11400'	27.7	14.0	50.5	14.0	14.3		
Crabtree Meadow	10700'	19.8	14.0	50.5 —	14.0	14.3		
Chagoopa Plateau	10700	21.8	5.0	22.9	5.0	7.0		
Pascoes	9150'	24.9	13.7	55.0	14.0	15.9		
Tunnel Guard Station	8900'	15.6	0.0	0.0	0.0	0.0		
Wet Meadows	8950'	30.3	0.0	0.0	0.0	0.0		
Casa Vieja Meadows	8300'	20.9	0.0	0.0	0.0	0.0		
Beach Meadows	7650'	11.0	0.0	0.0	0.0	0.0		
SURPRISE VALLEY AREA	. 000		0.0	0.0	0.0	0.0		
Dismal Swamp	7050'	29.2	26.9	92.1	26.4	28.7		
TRUCKEE RIVER								
Mount Rose Ski Area	8900'	38.5	36.9	95.8	36.3	38.1		
Independence Lake	8450'	41.4	45.3	109.4	45.3	44.8		
Big Meadows	8700'	25.7	10.1	39.3	10.0	11.0		
Squaw Valley	8200'	46.5	50.8	109.2	50.6	51.6		
Independence Camp	7000'	21.8	1.7	7.8	1.8	5.9		
Independence Creek	6500'	12.7	0.0	0.0	0.0	0.0		
Truckee 2	6400'	14.3	0.0	0.0	0.0	2.0		
LAKE TAHOE BASIN								
Heavenly Valley	8800'	28.1	11.5	40.9	11.5	12.5		
Hagans Meadow	8000'	16.5	0.8	4.8	0.9	0.9		
Marlette Lake	8000'	21.1	14.6	69.2	14.5	14.8		
Echo Peak 5	7800'	39.5	25.0	63.3	24.7	26.1		
Rubicon Peak 2 Tahoe City Cross	7500'	29.1 16.0	16.8	57.7 0.0	16.6	17.5		
Ward Creek 3	6750' 6750'	39.4	0.0 24.9	63.2	0.0 24.9	0.0 26.9		
Fallen Leaf Lake	6250'	7.0	0.0	0.0	0.0	0.0		
CARSON RIVER	0230	7.0	0.0	0.0	0.0	0.0		
Ebbetts Pass	8700'	38.8	26.3	67.8	26.0	27.4		
Poison Flat	7900'	16.2	0.0	0.0	0.0	0.0		
Monitor Pass	8350'	_	0.8	_	0.9	3.4		
Spratt Creek	6150'	4.5	0.0	0.0	0.0	0.0		
WALKER RIVER								
Leavitt Lake	9600'	_	54.8	_	54.6	54.4		
Virginia Lakes	9300'	20.3	12.4	61.1	12.4	12.2		
Lobdell Lake	9200'	17.3	4.9	28.3	4.9	6.9		
Sonora Pass Bridge	8750'	26.0	15.2	58.5	15.0	16.8		
Leavitt Meadows	7200'	8.0	0.0	0.0	0.0	0.0		
OWENS RIVER/MONO LAKE		a. =		ā : =				
Gem Pass	10750'	31.7	26.0	81.9	25.3	22.7		
Sawmill	10200'	19.4	4.9	25.2	4.9	7.5		
Cottonwood Lakes	10150'	11.6	0.0	0.0	0.0	0.0		
Big Pine Creek	9800'	17.9	1.8	10.3	1.8	3.1		
South Lake	9600'	16.0	0.0	0.0 71.6	0.0	2.9		
Mammoth Pass Rock Creek Lakes	9300' 10000'	42.4 14.0	30.4 0.0	71.6 0.0	30.0 0.0	28.9 0.0		
NOON CIEEN LANCS	10000	14.0	0.0	0.0	0.0	0.0		
NORMAL SNOWPACK ACCUMULATION EXPRESSED AS A PERCENT OF APRIL 1ST AVERAGE								

NORMAL SNOWPACK	ACCUMULATION	ON EXPRESSED AS	A PERCENT	OF APRIL 1ST	AVERAGE
AREA	JANUARY	FEBRUARY	MARCH	APRIL	MAY
Central Valley North	45%	70%	90%	100%	75%
Central Valley South	45%	4 ■ 65%	85%	100%	80%
North Coast	40%	15 ^{65%} _{60%}	85%	100%	80%





SNOWLINES

This year's annual meeting of the Western Snow Conference will be hosted by the South Continental Region. It will be held May 20–23, 2002 at Silver Creek Resort, a scenic 90 minute drive from Denver, CO. It is a little later this coming year and right before Memorial Day weekend. For further information regarding the Western Snow Conference contact Frank Gehrke at 916–574–2635 or gridley@water.ca.gov Information is available on the web at http://snobear.colorado.edu/WSC/WSC.html. Also appearing at this web site are last meeting's papers.

<u>Waiting</u> for runoff is a streamlogger in Budd Creek pictured on this months cover. Installed last summer as part of the intensive network of sensors in Yosemite National Park in a joint project involving Yosemite National Park, USGS, Scripps Institution of Oceanography/University of California, California Department of Water Resources and others.